

## **CERTIFICATE OF ANALYSIS**

Important Note:	Centrifuge before opening to ensure complete recovery of vial contents.		
Catalog #:	C01798M	Lot #:	49L33615
Description:	MAb to SEE Monoclonal Antibody to <i>Staphylococcus aureus</i> Enterotoxin E (SEE)		
Specificity:	Specific to <i>Staphylococcus aureus</i> enterotoxin E. No cross reaction to other types of <i>Staphylococcus aureus</i> enterotoxin A, B, C, and D.		
Clone:	B1918M		
Host Animal:	Mouse. Hybridization of P3X63.Ag8.653 myeloma cells with spleen cells from BALB/c mice.	Isotype:	IgG <sub>1</sub>
Immunogen:	Staphylococcus aureus Enterotoxin E		
Source:	Ascites		
Format:	Purified, Liquid		
Purification:	$\geq$ 95% pure (SDS-PAGE). Protein A Chromatography		
Concentration:	$3.2 \text{ mg/mL} (\text{OD280nm}, \text{E}^{0.1\%} = 1.38)$		
Buffer:	PBS: 1.5 mM Monopotassium Phosphate (KH <sub>2</sub> PO <sub>4</sub> ), 8.0 mM Disodium Phosphate (Na <sub>2</sub> HPO <sub>4</sub> ), 3.0 mM Potassium Chloride (KCl), 140 mM Sodium Chloride (NaCl), pH 7.4.		
Preservative:	0.05% Sodium Azide		
Applications:	Suitable for use in sandwich ELISA. Each laboratory should determine an optimum working titer for use in its particular application. Other applications have not been tested but use in such assays should not necessarily be excluded.Recommended antibody pair for sandwich immunoassay:Detection ControlCaptureDetection 		
Storage:	Short term (up to 7 days) store at 2-8°C. Long term, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles.		
Warning:	This product contains sodium azide, which has been classified as Xn (Harmful), in European Directive 67/548/EEC in the concentration range of $0.1 - 1.0$ %. When disposing of this reagent through lead or copper plumbing, flush with copious volumes of water to prevent azide build-up in drains.		

lobut att

Signature

03 Dec 2015 Date

FOR RESEARCH OR FURTHER MANUFACTURING USE ONLY