

<h2 style="margin: 0;">Aptamer Taq HS (Glycerol-Free), 50 U/μL</h2> <p style="margin: 10px 0 0 0;">For Research and Further Manufacturing use only</p>	Catalog No:	MDX015
	Lot No:	EN095-B127180
	Storage Conditions:	-20°C
	Component Lot No:	GFIH-324105A
	Expiry date:	June 2026

Quality Control Parameters

Analysis	Specification	Result
Functional	Activity is measured as DNA polymerase units by quantitative PCR analysis against a reference Taq DNA polymerase standard curve. <u>Pass Criteria:</u> Activity must be between 50 and 60 U/μL	59.30 U/μL
Functional	Aptamer specificity is	Passed
Glycerol content	Glycerol concentration is determined by spectrophotometric measurement of a colorimetric product from a coupled enzymatic reaction. <u>Pass Criteria:</u> Glycerol content <0.02 %	Passed
Purity	Purity is measured as a percentage of total protein by quantitative gel electrophoresis on Bioanalyzer (Agilent). <u>Pass Criteria:</u> >50 %	98.6 %
DNA contamination	DNA contamination is measured by quantitative PCR on <i>E. coli</i> and mouse genomic DNA specific targets. <u>Pass Criteria:</u> Amplification traces must overlay with the negative control.	Passed

United Kingdom

 Tel: +44 (0)20 8830 5300
 Fax: +44 (0)20 8452 2822

USA

 Tel: +1 901.382.8716
 Fax: +1 901.382.0027

Germany

 Tel: +49 (0)3371 60222 00
 Fax: +49 (0)3371 60222 01



Certificate of Analysis

COA No: CA_XBE-0068

Version: v04

DNase contamination	DNase contamination is measured as DNA substrate degradation against a DNase I dilution series by agarose gel electrophoresis. Limit of detection: 6.25×10^{-4} KU DNase I. <u>Pass Criteria:</u> No detectable degradation.	Passed
RNase contamination	RNase contamination is measured by quantitative PCR against RNase standards. Limit of detection: 9.7×10^{-3} ng/ μ L RNase. <u>Pass Criteria:</u> No detectable degradation.	Passed

QA / QC Representative:

J. Rahnenführer

Date: 08th May 2024

United Kingdom

Tel: +44 (0)20 8830 5300
Fax: +44 (0)20 8452 2822

USA

Tel: +1 901.382.8716
Fax: +1 901.382.0027

Germany

Tel: +49 (0)3371 60222 00
Fax: +49 (0)3371 60222 01

Enzyme Dilution Buffer

For research or further manufacturing use only


Catalog No:	MDX015
Lot No:	EN095-B127180
Storage Conditions:	-20°C
Lot number:	TDB-224105A
Expiry date:	June 2026

Quality Control Parameters

Enzyme Dilution Buffer is a glycerol-free 10x buffer used for the dilution of Taq antibody or reverse transcriptase.

Analysis	Specification	Result
Functional	<p>A 3Kb fragment was amplified with a dilution series of Enzyme Dilution Buffer, using standard conditions and 30 cycles.</p> <p>A 3Kb fragment was amplified with a dilution series of <i>Taq polymerase</i>, using standard conditions and 30 cycles.</p> <p>Single distinct bands were observed with agarose gel electrophoresis (ethidium stained).</p>	Passed
Glycerol determination	Glycerol content is < 0.2% determined by spectrophotometric analysis and comparison to a standard curve.	Passed
DNA contamination	Quantitative PCR analysis with no template. Presence of <i>E. coli</i> and mouse genomic DNA checked. Test sample must amplify in line with a reference sample.	Passed
DNase contamination	Incubation of a 1Kb double stranded DNA fragment. Incubation for 4 hours at 37°C with dilution series of DNase I. Analysed by agarose gel electrophoresis. Test sample must show less degradation than the limit of detection 6.25×10^{-4} KU/ μ L.	Passed
RNase contamination	Quantitative PCR analysis with high and low RNase standards. Test sample must show less RNase activity than the limit of detection: 9.7×10^{-3} ng/ μ L RNase.	Passed

QA / QC Representative:



J. Rahnenführer

 Date: 08th May 2024

United Kingdom

 Tel: +44 (0)20 8830 5300
 Fax: +44 (0)20 8452 2822

USA

 Tel: +1 901.382.8716
 Fax: +1 901.382.0027

Germany

 Tel: +49 (0)3371 60222 00
 Fax: +49 (0)3371 60222 01