

Lyo-Ready qPCR Mix

Product Handling Guide

Shipping: On Dry/Blue Ice
Catalog number: MDX021
Batch No.: See vial
Concentration: 2x

Store at -20 °C



Storage and stability:

Lyo-Ready qPCR Mix is shipped on dry/blue ice. On arrival store at -20 °C for optimum stability. Repeated freeze/thaw cycles should be avoided. Thawing during transportation does not affect the product performance. Solutions should be mixed/equilibrated after each thawing to avoid phasing.

Expiry:

When stored under the recommended conditions and handled correctly, full activity of the kit is retained until the expiry date on the outer box label.

Safety precautions:

Read and understand the SDS (Safety Data Sheets) before handling the reagents. Hardcopies of the SDS will be provided with the first shipment, thereafter they will be available upon request.

Quality control:

Bioline operates under ISO 13485 Quality Management System. Lyo-Ready qPCR Mix and its components are extensively tested for activity, processivity, efficiency, heat activation sensitivity, absence of nuclease contamination and absence of nucleic acid contamination.

Notes:

This reagent has been manufactured under 13485 Quality Management System and is suitable for further manufacturing use as an IVD component.

Description

Lyo-Ready qPCR Mix is a glycerol-free qPCR mix containing Taq polymerase, reaction buffer, dNTP, MgCl₂ and lyo-exipients. In order to produce room temperature lyophilized qPCR reagents, assay specific primers and probes are added to Lyo-Ready qPCR Mix for subsequent lyophilization.

Kit components

Table 1

Component
Lyo-Ready qPCR Mix, 2x

Users Guidelines

Master mix preparation

Recommended reagent volumes per 20 µL qPCR mix are given in Table 2.

Table 2

Reagent	Volume
Lyo-Ready qPCR Mix, 2x	10 µL
Primer-Probe Mix, 20x	1 µL
Water	x µL
Total volume	Up to 20 µL

Dispense into reaction vials and immediately transfer into a freeze-dryer and run a suitable drying cycle.

For long-term storage at ambient temperatures, the lyophilized product from the freeze-dryer should be packaged at low relative humidity conditions.

Assay setup

Rehydrate the lyophilized qPCR mix in the reaction vials with 20 µL template-containing solution and run qPCR.

The qPCR conditions in Table 3 are suitable for amplicons of up to 200 bp. These cycling parameters have been optimized for Lyo-Ready qPCR Mix on a number of platforms, however they can be varied to suit different machine-specific protocols.

Table 3

Step	Temperature	Time	Cycles
Polymerase activation	95 °C	2 min	1
Denaturation	95 °C	5 s	45
Annealing/Extension	60 °C	20 s	

Technical Support

For any technical enquiries, please contact our Technical Support team via email at: mbi.tech@meridianlifescience.com

Bioline Reagents Ltd
UNITED KINGDOM

Tel: +44 (0)20 8830 5300
Fax: +44 (0)20 8452 2822

Bioline USA Inc.
USA

Tel: +1 901 382 8716
Fax: +1 901 382 0027

Bioline GmbH
GERMANY

Tel: +49 (0)337 168 1229
Fax: +49 (0)3371 68 1244

Bioline (Aust) Pty. Ltd
AUSTRALIA

Tel: +61 (0)2 9209 4180
Fax: +61 (0)2 9209 4763

Bioline France
FRANCE

Tel: +33 (0)1 42 56 04 40
Fax: +33 (0)9 70 06 62 10