

Inhibitor-Tolerant RT-qPCR Lo-ROX Mix

Product Handling Guide

Shipping:	On dry/blue ice
Catalog number:	MDX105
Batch No.:	See vial
Concentration:	4x

Store at -20 °C



Storage and stability:

Inhibitor-Tolerant RT-qPCR Lo-ROX Mix is shipped on dry/blue ice. On arrival store at -20 °C for optimum stability. Repeated freeze/thaw cycles should be avoided. Thawing during transportation does not affect the product performance. Solutions should be mixed/equilibrated after each thawing to avoid phasing.

Expiry:

When stored under the recommended conditions and handled correctly, full activity of the kit is retained until the expiry date on the outer box label.

Safety precautions:

Read and understand the SDS (Safety Data Sheets) before handling the reagents. Hardcopies of the SDS will be provided with the first shipment, thereafter they will be available upon request.

Quality control:

Meridian operates under ISO 13485 Quality Management System. Inhibitor-Tolerant RT-qPCR Lo-ROX Mix and its components are extensively tested for activity, processivity, efficiency, heat activation sensitivity, absence of nuclease contamination and absence of nucleic acid contamination

Notes:

For research and/or further manufactured use only.

Description

Inhibitor-Tolerant RT-qPCR Lo-ROX Mix is a one tube formulation combining the latest advances in buffer chemistry and PCR enhancers, together with an optimized concentration of antibody-mediated hot-start polymerase, reverse transcriptase, RNase Inhibitors, dNTPs and MgCl₂. Inhibitor-Tolerant RT-qPCR Mix has been designed for highly reproducible, accurate RNA and DNA target amplification under fast thermal cycling conditions, delivering excellent results in multiplex assays, even in the presence of difficult inhibitors found in sputum and stool samples.

Kit components

Table 1

Component
Inhibitor-Tolerant RT-qPCR Lo-ROX, 4x

Users Guidelines

Master mix preparation

Recommended reagent volumes per 20 µL qPCR mix are given in Table 2.

Table 2

Reagent	Volume
Inhibitor-Tolerant RT-qPCR Lo-ROX Mix, 4x	5 µL
Primer-Probe Mix, 20x	1 µL
Template*	Up to 10 µL
Water	As required
Total volume	Up to 20 µL

* RNA and DNA template from extracted or crude sample lysate. Template volume optimization required.

Assay setup

The qPCR conditions in Table 3 are suitable for amplicons of up to 200 bp. These cycling parameters have been optimized for Inhibitor-Tolerant RT-qPCR Lo-ROX Mix on a number of platforms, however they can be varied to suit different machine-specific protocols.

Table 3

Step	Temperature	Time	Cycles
Reverse transcription**	50 °C	10 min	1
Polymerase activation	95 °C	2 min	1
Denaturation	95 °C	5 s	45
Annealing/Extension**	60 °C	20 s	

**When multiplexing, the reverse transcription reaction time can be extended up to 20 minutes and/or the temperature can be increased up to 55° C and the annealing/extension time can be extended up to 60 seconds and/or the temperature can be increased up to 65° C.

Related Products	Cat. No.
Inhibitor-Tolerant qPCR Mix	MDX013
Inhibitor-Tolerant RT-qPCR Mix	MDX016
Fast 1-Step RT-qPCR Mix	MDX018
Lyo-Ready 1-Step RT-qPCR Mix	MDX024

Technical Support

For any technical enquiries, please contact our Technical Support team via email at: mbi.tech@meridianlifescience.com